LaTeX to PDF and MathJax: Example 1

Emma Cliffe

2017

Contents

| Using this document | | | | | |
|---------------------|-----|-----------------------------------|---|--|--|
| 1 | Qua | dratic equations | 2 | | |
| | 1.1 | Solutions to a quadratic equation | 2 | | |
| | 1.2 | The discriminant | 2 | | |

List of Figures

| 1 Exa | mples of quadra | ic functions with zer | o, one and two real roots. | | 3 |
|-------|-----------------|-----------------------|----------------------------|--|---|
|-------|-----------------|-----------------------|----------------------------|--|---|

List of Tables

1 Number of real roots of a quadratic equation, given the discriminant . . 3

Using this document

This is an example of a document compiled from LATEX into multiple formats:

- Standard print PDF
- Clearer print PDF
- Accessible web format
- Accessible Word document

The outputs can be used to test setups and as a first example for students to try out.

1 Quadratic equations

A quadratic equation is an equation with the form $ax^2 + bx + c = 0$ where x represents an unknown and a, b and c are known numbers with $a \neq 0$.

1.1 Solutions to a quadratic equation

A solution to a quadratic equation is a value of x such that the equation balances. The solutions to quadratic equations can be found by using the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$
(1)

Example.

For instance, the solutions to $x^2 + 2x - 3 = 0$ are:

$$x = \frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times -3}}{2 \times 1}$$
$$= \frac{-2 \pm \sqrt{4 + 12}}{2}$$
$$= \frac{-2 \pm \sqrt{16}}{2}$$
$$= \frac{-2 \pm 4}{2}$$

Hence, x = 1 or x = -3.

1.2 The discriminant

Definition (Discriminant).

The **discriminant** of a quadratic equation with coefficients $a, b, c \in \mathbb{R}$ is:

$$\Delta = b^2 - 4ac.$$

Remark.

Note that this is the expression beneath the square root symbol in the quadratic formula (1).

We can use the discriminant to determine the number of real roots of a quadratic equation. The number depends on the value of Δ as in table 1.

| Value of Δ | Real roots |
|-------------------|---------------|
| $\Delta > 0$ | Two, distinct |
| $\Delta = 0$ | One, repeated |
| $\Delta < 0$ | Zero |

Table 1: Number of real roots of a quadratic equation, given the discriminant

Figure 1 shows an example of each possibility¹.




Figure 1: Examples of quadratic functions with zero, one and two real roots.

¹The image is due to Olin, CC-BY-AS 3.0 downloaded from Wikimedia Commons